

TREATMENT OF BACTERIAL MDR (MRSA, PRSA, VRE) WITH BOTANICAL THERAPIES

David Winston, RH (AHG)

References

- Abascal K and E Yarnell. Herbs and drug resistance, part I – herbs and microbial resistance to antibiotics. *Alt Comp Ther* 2002; 8(4):237-241.
- Abascal K and E Yarnell. Herbs and drug resistance, part II – herbs and microbial resistance to antibiotics. *Alt Comp Ther* 2002; 8(5):284-290.
- Aqil F et al. Effect of certain bioactive plant extracts on clinical isolates of beta-lactamase producing methicillin resistant *Staphylococcus aureus*. *J Basic Microbiol* 2005; 45(2):106-14.
- Ahmed I and A Beg. Antimicrobial and phytochemical studies on 45 Indian medicinal plants against multi-drug resistant human pathogens. *J Ethnopharm* 2001; 74(2):113-23.
- Anonymous. Invasive MRSA. *CDC Fact Sheet* 2007; http://www.cdc.gov/ncidod/dhqp/ar_mrsa_invasive_FS.html.
- Anonymous. Honey makes medical comeback. *Associated Press*, 2007.
- Anonymous. Staph germ undermines body's defenses. *Associated Press*, 2007.
- Cegiela-Carlio P et al. Modulation of multi-drug resistance (MDR) in *Staphylococcus aureus* by Osha (*Ligusticum porteri* L. Apiaceae) essential oil compounds. *Flav Frag J* 2005; 20(6):671-5.
- Chen L et al. Inhibition of growth of *Streptococcus mutans*, methicillin-resistant *Staphylococcus aureus*, and vancomycin-resistant enterococci by kurarinone, a bioactive flavonoid isolated from *Sophora flavescens*. *J Clin Microbiol* 2005; 43(7):3574-5.
- Dryden M et al. A randomized, controlled trial of tea tree topical preparations versus a standard topical regimen for the clearance of MRSA colonization. *J Hosp Infect* 2004; 56(4):283-6.
- Freestone P et al. Microbial endocrinology: how stress influences susceptibility to infection. *Trends in Microbiology* 2008; 16(2):55-64.
- Gibbons S et al. Activity of *Zanthoxylum clava-herculis* extracts against multi-drug resistant methicillin-resistant *Staphylococcus aureus* (mdr-MRSA). *Phytother Res* 2003; 17(3):274-5.
- Gibbons S et al. Bacterial resistance modifying agents from *Lycopus europaeus*. *Phytochemistry* 2003; 62(1):83-7.
- Guz N et al. Flavonolignan and flavone inhibitors of a *Staphylococcus aureus* multidrug resistance pump: structure-activity relationships. *J Med Chem* 2001; 44(2):261-68.
- Harris J et al. Antimicrobial properties of *Allium sativum* (garlic). *Appl Microbiol Biotechnol* 2001; 57(3):282-6.
- Hatano T et al. Phenolic constituents of *Cassia* seeds and antibacterial effect of some naphthalenes and anthraquinones on methicillin-resistant *Staphylococcus aureus*. *Chem Pharm Bull (Tokyo)* 1999; 47(8):1121-7.

- Hatano T et al. Phenolic constituents of licorice. VIII. Structures of glicophenone and glicoisoflavanone, and effects of licorice phenolics on methicillin-resistant *Staphylococcus aureus*. *Chem Pharm Bull (Tokyo)* 2000; 48(9):1286-92.
- Haydel S and L Williams. Broad-spectrum *in vitro* antibacterial activities of clay minerals against antibiotic-susceptible and antibiotic-resistant bacterial pathogens. *Geological Society of America Annual Meeting*, 2007.
- Horiuchi K et al. Potentiation of antimicrobial activity of aminoglycosides by carnosol from *Salvia officinalis*. *Biol Pharm Bull* 2007; 30(2):287-90.
- Huycke M et al. Multiple-drug resistant enterococci: the nature of the problem and an agenda for the future. *Emerg Infect Dis* 1998; 4(2):239-49.
- Kamijo M, F Kanazawa, and M Funaki. Effects of *Rosa rugosa* petals on intestinal bacteria. *Bioscience, Biotechnology, Biochemistry* 2008; 706450I-5.
- Kassam M et al. Green tea shows superbug-battling potential. www.nutraingredients-usa.com/new/printNewsBis.asp?id=84356.a
- Kim K et al. Antibacterial activity of *Curcuma longa* L. against methicillin-resistant *Staphylococcus aureus*. *Phytother Res* 2005; 19(7):599-604.
- Klevins R et al. Invasive methicillin-resistant *Staphylococcus aureus* infections in the United States. *JAMA* 2007; 298(15):1763-71.
- LaPlante K. *In vitro* activity of lysostaphin, mupirocin, and tea tree oil against clinical methicillin-resistant *Staphylococcus aureus*. *Diagn Microbiol Infect Dis* 2007; 57(4):413-8.
- Lechner D et al. The anti-staphylococcal activity of *Angelica dahurica* (Bai Zhi). *Phytochemistry* 2004; 65(3):331-5.
- Liu I and D Durhan. Baicalin synergy with beta-lactam antibiotics against methicillin-resistant *Staphylococcus aureus* and other beta-lactam-resistant strains of *S. aureus*. *J Pharm Pharmacol* 2000; 52(3):361-6.
- Liu M and S Matsuzak. Antibacterial activity of flavonoids against methicillin-resistant *Staphylococcus aureus* (MRSA). *Dokkyo J Med Sci* 1995; 22:253-61.
- Lugman S et al. Potential of rosemary oil to be used in drug resistant infections. *Altern Ther Health Med* 2007; 13(5):54-9.
- Natarajan, P., Katta, S., et al, Positive Antibacterial Co-Action Between Hop (*Humulus lupulus*) Constituents and Selected Antibiotics, *Phytomedicine*, 15 (2008):194-201
- Nayak B et al. The evidence based wound healing activity of *Lawsonia inermis* Linn. *Phytother Res* 2007; 21(9):827-31.
- Nishida S. Effect of Hochu-ekki-to on asymptomatic MRSA bacteriuria. *J Infect Chemother* 2003; 9:58-61.
- Nostro A and A Cannatelli. The effect of *Nepeta cataria* extract on adherence and enzyme production of *Staphylococcus aureus*. *Int J Antimicrob Agents* 2001; 18(6):583-5.
- Oluwatuyi M et al. Antibacterial and resistance modifying activity of *Rosmarinus officinalis*. *Phytochemistry* 2004; 65(24):3249-54.

- Onlen Y et al. Antibacterial activity of propolis against MRSA and synergism with topical mupirocin. *J Alt Comp Med* 2007; 13(7):713-8.
- Quinlan A. MRSA faces defeat from wild flower. *CIT*, 2007.
- Rouveix B. Clinical implications of multiple drug resistance efflux pumps of pathogenic bacteria. *J Antimicrob Chemother* 2007; 59(6):1208-9.
- Rudrappa T and H Bais. Curcumin, a known phenolic from *Curcuma longa*, attenuates the virulence of *Pseudomonas aeruginosa* PA01 in whole plant and animal pathogenicity models. *Journal of Agricultural and Food Chemistry* 2008; 56:1955-62.
- Shiota S et al. Mechanisms of action of corilagin and tellimagrandin I that remarkably potentiate the activity of beta-lactams against methicillin-resistant *Staphylococcus aureus*. *Microbiol Immunol* 2004; 48(1):67-73.
- Smith E et al. Isopimaric acid from *Pinus nigra* shows activity against multidrug-resistant and EMRSA strains of *Staphylococcus aureus*. *Phytother Res* 2005; 19(6):538-42.
- Song J et al. *In vitro* inhibitory effects of *Polygonum cuspidatum* on bacterial viability and virulence factors of *Streptococcus mutans* and *Streptococcus sobrinus*. *Arch Oral Biol*, 2006, Dec;51(12):1131-40.
- Stavri M et al. Bacterial efflux pump inhibitors from natural sources. *J Antimicrob Chemother* 2007; 59:1247-60.
- Stermitz F et al. 5'-methoxyhydnocarpin-D and pheophorbide A: *Berberis* species components that potentiate berberine growth inhibition of resistant *Staphylococcus aureus*. *J Nat Prod* 2000; 63(8):1146-49.
- Tegos G et al. Multidrug pump inhibitors uncover remarkable activity of plant antimicrobials. *Antimicrob Agents Chemother* 2002; 46(10):3133-41.
- Voravuthikunchai S and L Kitpipit. Antibacterial activity of crude extracts of Thai medicinal plants against clinical isolates of methicillin-resistant *Staphylococcus aureus*. *Songklanakarin J Sci Technol* 2005; 27:525-34.
- Weckesser S et al. Screening of plant extracts for antimicrobial activity against bacteria and yeasts with dermatological relevance. *Phytomedicine*. 2007; 14(7-8):508-16.
- Williams S. Clay that kills: ground yields antibacterial agents. *Science News* 2007; 172(18):276.
<http://www.sciencenews.org/articles/20071103/fob4.asp>.
- Winston D and S Maimes. *Adaptogens: Herbs for Strength, Stamina, and Stress Relief*. Healing Arts Press: Rochester, VT, 2007.
- Yang Z et al. The synergistic activity of antibiotics combined with eight traditional Chinese medicines against two different strains of *Staphylococcus aureus*. *Colloids Surf B Biointerfaces*. 2005; 41(2-3):79-91.
- Yarnell E and K Abascal. The complicated web of toxins, drug resistance, and herbs. *Alt Comp Ther* 2003; 9(1):16-18.
- Yu H et al. Antimicrobial activity of berberine alone and in combination with ampicillin or oxacillin against methicillin-resistant *Staphylococcus aureus*. *J Med Food* 2005; 8(4):454-61.
- Zaidan M et al. *In vitro* screening of five local medicinal plants for antibacterial activity using disc diffusion methods. *Trop Biomed*. 2005; 22(2):165-70.